

Foreword

Although adverse health effects of exposures to high level of arsenic and/or fluoride through drinking water originated from underground sources are not new in China or other regions, it is now recognized as one of the global environmental issues. It has been revealed that more than 100 million people in Bangladesh, West Bengal in India, China etc. are exposed to toxic dose of arsenic, among whom very serious cases of not only skin symptoms/lesions but also various cancers of skin, lung etc. have been occurring. (in case of fluoride intoxication its typical effect is called as fluorosis. Its combination effects with arsenic have also been suggested) It should be noted that air pollution with arsenic and/or fluoride through coal-burning may be another source of exposures related to their intoxications.

On the other hand, this issue is not only the problem for those developing countries. The developed societies have also become inescapable from those serious environmental pollutions, traditional type or modern type, in the developing countries through their transboundary transports accelerated with rapid internationalization of world trades, migrations etc., indicating that more emphasis should be given regarding such risk managements at the international level.

In every sense, I hope, on behalf of the participants to this workshop, this publication would be meaningful in promoting and extending studies and managements related to this one of the most serious environmental issues in the present world.

Michinori KABUTO, DHSc

Acting Director for Health Researches
National Institute for Environmental Studies
16-2 Onogawa, Tsukuba, Ibaraki,
305-8506 JAPAN
tel:+81-298-50-2333 fax:+81-298-50-2571
e-mail: kabuto@nies.go.jp

Preface

This workshop was planned as part of an HDP (Human Dimension Programme for global environmental issues) study on arsenic pollution due to increasing uses of underground water and environmental planning, which was purposed to evaluate and summarize the state of knowledge and findings and also to consider the appropriate mitigation or alternative methods to stop as early as possible the expected huge amount of health burdens. To this workshop, 5 scientists were invited from India, Thailand and China to provide an overview of the general situations of arsenism and researches, and also two Japanese scientists who have been working on Bangladesh and China (Inner Mongolia), respectively for a brief introduction of their own researches as well as general comments and recommendations.

The time schedule was as follows:

10:00-10:15	Opening remarks	Gen Ohi (Director, NIES)
10:15-15:00	Current Status of Arsenic Pollution and Health Impacts in	
10:15-12:00	West Bengal, Bangladesh and Thailand	
	Arsenic pollution in West-Bengal India and Bangladesh	
	Dipankar Chakraborti (School of Environmental Studies, Jadavpur University)	
	Arsenicism in Ronphibun, Thailand~ past, present and future	
	Shouko Oshikawa (Epidemiology Unit, Prince of Songkla University, Thailand)	
	Comments and Discussion	
	An arsenism study on communities in Bangladesh	
	Chiho Watanabe (School of International Health, University of Tokyo)	
12:00-13:00	Lunch	
13:00-15:00	China	
	Progress on Arsenism Research in China	
	Jin Yinlong (IEHE, Chinese Academy of Preventive Medicine)	
	Current Situation of Arsenism in China	
	Sun Guifan (School of Public Health, China Medical University)	
	Assessment of Arsenic Removal from Drinking Water by New Adsorbents	
	He Gongli (IEHE, Chinese Academy of Preventive Medicine)	
15:00-15:30	Coffee Break	
15:30-16:15	Comments and Discussion	
	Study of arsenism via consumption of arsenic contaminated ground water and intervention study in water mitigation in Inner Mongolia	
	Takahiko Yoshida (Asahikawa Medical College)	
16:15-17:00	Q&A and Discussion Michinori Kabuto (NIES)	
17:00	Adjournment	

Contents

Foreword

Preface

Session 1: Current Status of Arsenic Pollution and Health Impacts in West Bengal, Bangladesh and Thailand

- 1.1 Arsenic pollution in West-Bengal India and Bangladesh
Dipankan Chakraborti ····· 1
- 1.2 Arsenicism in Ronphibun, Thailand- past, present and future
Shoko Oshikawa ····· 16
- 1.3 Comments and Discussion
Chiho Watanabe ····· 26

Session 2: Current Status of Arsenic Pollution and Health Impacts in China

- 2.1 Current situation of arsenism in China
Sun Guifan ····· 31
- 2.2 Progress on the research of arsenism in China
Jin Yinlong and Li You ··· 34
- 2.3 Assessment of arsenic Removal from drinking water by new adsorbents
He Gongli ····· 40
- 2.4 Comments and Discussion
Takahiko Yoshida ····· 45

Session 3: Concluding Remarks

- Major Points Confirmed and Recommendations for Future Studies
Michinori Kabuto ····· 55
- Participants List ····· 57

Session 1

Current Status of Arsenic Pollution and Health Impacts in West Bengal, Bangladesh and Thailand

1.1 Current Status of Arsenic Pollution and Health Impacts in West Bengal and Bangladesh

Mohammad Mahmudur Rahman¹, Kunal Paul¹, Uttam K. Chowdhury¹, Bhajan K. Biswas¹, Dilip Lodh¹, Gautam K. Basu¹, Shibtosh Roy², Ranajit Das², Bashir Ahmed², Imrul Kaies², Ajoy Kishore Barua², Shyamal K. Palit², Quazi Quamruzzaman² and Dipankar Chakraborti^{1*}

1 School of Environmental Studies, Jadavpur University, Calcutta-700032, India

2 Dhaka Community Hospital, Dhaka - 1217, Bangladesh

School of Environmental Studies (SOES), Jadavpur University, Calcutta, India is working on arsenic groundwater contamination in West Bengal-India for last 13 years and for last 6 years in Bangladesh jointly with Dhaka Community Hospital (DCH), Bangladesh. **Table-1** shows the overall arsenic situation in West Bengal-India ; **Table-1(A)** detailed study of one out of the nine affected districts; **Table- 1(B)** one out of the 74 affected Blocks and **Table-1(C)** one out of 2000 affected villages. **Table-2** shows the arsenic situation in Bangladesh; **Table-2(A)** detailed study of one out of 47 affected districts; **Table-2(B)** one out of 178 affected Police Stations and **Table-2(C)** one out of 2500 affected villages.

The overall arsenic groundwater contamination in West Bengal-India and Bangladesh is serious. More than 100 million people are living in arsenic affected districts. That does not mean all are drinking arsenic contaminated water but no doubt population are at risk. Although we are surveying for last 13 years and 6 years in arsenic affected districts of West Bengal and Bangladesh with about 40 people in our group but still we feel we have seen the tip of the iceberg. More we are surveying more and more people with arsenical skin lesions and affected areas are coming to limelight.

Arsenical skin lesion is late manifestation. When skin-lesions will appear on body depends on many factors like (a) concentration of arsenic in drinking water (b) how long one is drinking the contaminated water (c) how much one is drinking contaminated water per day (in many families we have found those drinking too much water have arsenic skin lesions while those drinking less have no skin lesions) (d) nutrition status of the people drinking contaminated water also plays an important roles. We have hundreds of example showing that poor people suffering from malnutrition showing arsenical skin lesions but people getting better nutrition are not showing any skin lesions drinking the same concentration (also close to same volume) of contaminated water. From our 13 years field survey screening more than 100,000 people from West Bengal and Bangladesh and having a list of 12000 registered arsenic

patients with skin lesions, we can say that usually above 300 µg/l of arsenic in drinking water may show arsenical skin lesions (exceptions are always there). However, we have found a few patients with spotted melanosis drinking 90 µg/l (Kolsur village, Deganga Block, North 24-Parganas) of arsenic in drinking water in West Bengal and 80 µg/l in Samta village, Sarsa Police Station of Jessore District of Bangladesh. However these are exceptions. Children below 11 years normally do not show the skin lesions but we have registered a few hundred patients aged below 11 years from West Bengal, India and Bangladesh, when arsenic in their drinking water is quite high around 700µg/l and above. We have also observed that children suffering from malnutrition may also suffer from arsenical skin lesions even with lower arsenic in their drinking water (around 400 µg/l).

We have also found good co-relation between arsenic in hair, nail, urine and concentration of arsenic in drinking water.

After analyzing 32960 water samples from hand tube-wells of Bangladesh, we had noticed that in some parts of Bangladesh, the arsenic contamination in groundwater is minimum, some parts are almost arsenic contamination free and some parts are highly contaminated. While trying to find out

* Corresponding author; Tel: 91 33 4735233, Fax: 91 33 4734266, Email: dcsoesju@vsnl.com

the reason we noticed that out of the four geo-morphological regions of Bangladesh (a) Hill Tract (b) Flood Plain (c) Pleistocene Plain and Pleistocene Upland and (d) Deltaic Plain and Coastal Plain, Hill tract and Pleistocene Plain and Pleistocene Upland are usually arsenic contamination free. Some Flood Plain areas between Pleistocene Plain and Pleistocene Upland like Nilphamari, Lalmonirhat, Kurigram, Rangpur districts are also less contaminated. We have further noticed that in fringe area of Pleistocene Plain and Pleistocene Upland with Flood Plain; Hill Tract with Flood Plain and if river had eroded Pleistocene Plain and Pleistocene Upland and Hill Tract areas, some contamination are there (the probable reason may be heavy deposition of Holocene sediments). Deltaic Plain with Coastal belt and Flood Plain regions are the most arsenic contamination area of Bangladesh. At present about 44.2% of the water samples are above 50 µg/l in Deltaic Plain with Coastal belt and Flood Plain but from our West Bengal experience we expect in future more tube-wells in these areas may get arsenic contamination.

While analyzing of deep tube wells above 100 meter in 4 geo-morphological regions of Bangladesh, we found all tube wells above 100m in Pleistocene Plain and Pleistocene Upland and Hill tract are also arsenic contamination free (< 10 µg/l). In case

of flood plain and Deltaic plain (including coastal plain) we had noticed that deep tube wells above 300m are mostly safe with respect to 50 µg/l but about 22% of tube wells above 300 m contain arsenic between 10 and 50 µg/l.

The source of arsenic in West Bengal-India and Bangladesh is geogenic. We have identified arsenic rich pyrite in bore-hole sediment samples.

Table-1: Overall arsenic situation in West Bengal-India

	West Bengal-India
Area in sq. km	89193
Population in million (according to 1991 census)	68
Total number of districts	18
Number of arsenic affected districts (groundwater arsenic above 50 µg/l)	9
Total number of water samples analyzed	100000
% of samples having arsenic >10µg/l	51
% of samples having arsenic > 50µg/l	34
Area of arsenic affected districts in sq. km	38865
Population of arsenic affected districts in million	42.7
Number of arsenic affected Blocks / Police Stations	74
Number of arsenic affected villages (approx.) where groundwater >50 µg/l	2000
People drinking arsenic contaminated water >50 µg/l (in million)	6
Districts surveyed for arsenic patients	7
Patients found from surveyed districts	7
People screened for arsenic patients from affected villages (preliminary survey)	86000
Number of registered patients with clinical manifestations	8500 (9.8%)
% of children having arsenical skin lesions of total patients	1.7
Total hair, nail, urine, skin-scales analyzed	27000
Arsenic above normal level in biological samples	77%

Table-1A: Arsenic situation in North 24-Pargana one of the nine arsenic affected districts of West Bengal-India

Area of the district in sq. km	4094
Population of the district in million	7.3
Number of Blocks / Police Stations in North 24-Parganas	22
Number of blocks / Police Stations surveyed	22
Number of hand tubewell's water sample analyzed	28872

Number of samples having arsenic >10 µg/l	15513 (53.7%)
Number of samples having arsenic >50 µg/l	9339 (32.3%)
Total blocks having arsenic >10 µg/l	20
Total blocks having arsenic >50 µg/l	19
Total blocks where arsenic patients identified	16
Total hair, nail, urine, skin-scales analyzed	21158
Arsenic above normal level (average) in biological samples	74.3%
Total population screened for arsenic patients from affected villages (preliminary survey)	15902
Number of registered patients with clinical manifestations	1622 (10.2%)
People drinking arsenic contaminated water >50 µg/l in million	1.15

Table-1B: Arsenic situation in Deganga, North 24-Paraganas one block out of 74 affected blocks in West Bengal-India

Area of the Block in sq. km	201
Population of the Block (according to 1991 census)	234141
Total hand tubewells in Deganga block	15886
Number of hand tubewell's water sample analyzed	8785 (55.3%)
Number of samples having arsenic >10 µg/l	5082 (57.8%)
Number of samples having arsenic >50 µg/l	3610 (41.1%)
Total hair, nail, urine, skin-scales analyzed	17365
Arsenic above normal level (average) in biological samples	67.6%
Total population screened for arsenic patients in affected villages (preliminary survey)	1574
Number of registered patients with clinical manifestations	409 (26%)
People drinking arsenic contaminated water >50 µg/l	96209
Number of hand tubewell's water analyzed from schools	127
Number of samples having arsenic >10 µg/l	74 (58.2%)
Number of samples having arsenic >50 µg/l	62 (48.8%)
Amount of arsenic falling on agricultural land from 3200 big-diameter irrigation tubewells in one year	6.4 tons

Table-1C: Arsenic situation in Kolsur village of Deganga block, North 24-Parganas one of 2000 affected villages of West Bengal-India

Area of Kolsur in sq. km	16.32
Population of Kolsur	16,879
Total number of hand tubewells in Kolsur	1550
Number of hand tubewell's water sample analyzed	1184
Number of samples having arsenic >10 µg/l	961 (81.1%)

Number of samples having arsenic >50 µg/l	859 (72.5%)
Total hair, nail, urine, skin-scales analyzed	3633
Arsenic above normal level (average) in biological samples	93.3%
Total population screened for arsenic patients (detailed survey)	11000
Number of registered patients with clinical manifestations	579 (5.27%)
People drinking arsenic contaminated water >50 µg/l	12238

Table-2: Overall arsenic situation in Bangladesh

	Bangladesh
Area in sq. km	148393
Population in million	120
Total number of districts	64
Total number of water samples analyzed	32960
% of samples having arsenic >10µg/l	56.3
% of samples having arsenic > 50µg/l	37.3
Number of arsenic affected districts (groundwater arsenic above 50 µg/l)	47
Area of arsenic affected districts in sq. km	112407
Population of arsenic affected districts in million	93.4
Number of arsenic affected Blocks / Police Stations	178
Number of arsenic affected villages (approx.) where groundwater >50 µg/l	2500
People drinking arsenic contaminated water >50 µg/l (in million)	25
Districts surveyed for arsenic patients	34
Patients found from surveyed districts	32
People screened for arsenic patients (preliminary survey)	17896
Number of registered patients with clinical manifestations	3695 (20.6%)
% of children having arsenical skin lesions of total patients	6.4
Total hair, nail, urine, skin-scales analyzed	10496
Arsenic above normal level (average) biological samples	93%

Table-2 A: Arsenic situation in district Lakshmipur one of the 47 arsenic affected districts of Bangladesh

Area of the district in sq. km	1456
Population of the district in million	1.5
Number of Blocks / Police Stations in Lakshmipur	4
Number of blocks / Police Stations surveyed	4
Number of hand tubewell's water sample analyzed	2662

Number of samples having arsenic >10 µg/l	2358 (88.6%)
Number of samples having arsenic >50 µg/l	2123 (79.8%)
Number of Police Stations having arsenic >10 µg/l	4
Number of Police Stations having arsenic >50 µg/l	4
Number of Police Stations surveyed for arsenic patient	3
Number of Police Stations where arsenic patients identified	3
Number of villages surveyed for arsenic patient	27
Number of villages where arsenic patients identified	25
Number of people screened for arsenical skin lesions (preliminary survey)	2138
Number of registered patients with clinical manifestations	504 (23.2%)
Number of the children (<11 years) screened for arsenical skin lesions	400
Number of registered child patients with clinical manifestations	25 (6.2%)
People drinking arsenical contaminated water >50 µg/l in million	1.2 (85%)
Total hair, nail, urine, skin-scales analyzed	1250
Arsenic above normal level (average) in biological samples	92%

Table-2B: Arsenic situation in Police Station Ramganj in Lakshmipur district out of 174 affected Police Stations in Bangladesh

Area of Ramganj in sq. km	169
Population of Ramganj	274000
Number of hand tubewell's water sample analyzed	2077
Number of samples having arsenic >10 µg/l	1853 (89%)
Number of samples having arsenic >50 µg/l	1668 (80.3%)
Total hair, nail, urine, skin-scales analyzed	503
Arsenic above normal level (average) in biological samples	92%
Total population screened for arsenic patients (preliminary survey)	970
Number of registered patients with clinical manifestations	300 (31%)
Number of children (<11 years) screened for arsenical skin lesions	183
Number of registered child patients with clinical manifestations	11 (6%)
People drinking arsenic contaminated water >50 µg/l	164390

Table-2C: Arsenic situation in one village Samta of Sarsa Police Station in Jessore district of Bangladesh

Area of Samta in sq. km	3.2
Population of Samta	4841
Total number of hand tubewells in Samta	276
Number of hand tubewell's water sample analyzed	265
Number of samples having arsenic >10 µg/l	260 (98.1%)

Number of samples having arsenic >50 µg/l	242 (91.3%)
Total hair, nail, urine, skin-scales analyzed	830
Arsenic above normal level (average) in biological samples	96.3%
Total population screened for arsenic patients (detailed survey)	600
Number of registered patents with clinical manifestations	330 (55%)
People drinking arsenic contaminated water >50 µg/l	4356

List of publications of School of Environmental Studies on arsenic groundwater contamination in West Bengal-India and Bangladesh

1992

Environmental Pollution & Chronic Arsenicosis in South Calcutta, West Bengal. D.N.Guha Mazumder, J.Das Gupta, A.K. Chakraborty, A. Chatterjee, D.Das & D.Chakraborti; Bulletin of World Health Organization; 70 (4), 481-485, 1992.

1993

A Study of Ground Water Contamination by Arsenic in the Residential Area of Behala, Calcutta due to Industrial Pollution. Amit Chatterjee, Dipankar Das & D.Chakraborti; Environmental Pollution, 80 (1), 57-65, 1993.

1994

Arsenic Contamination in Ground Water in Six Districts of West Bengal: the Biggest Arsenic Calamity in the World. D. Das, A. Chatterjee, G.Samanta, B. Mandal, T. Roy Chowdhury, G. Samanta, P. P. Chowdhury, C. Chanda, G. Basu, D. Lodh, S. Nandi, T. Chakraborty, S. Mandal, S. M. Bhattacharya and D.Chakraborti, Analyst, 119, 168N-170N, 1994.

1995

Arsenic in Ground Water in Six Districts of West Bengal, India: The Biggest Arsenic Calamity in the World Part 1. Arsenic Species in Drinking Water and Urine of the Affected People. Amit Chatterjee, Dipankar Das, Badal K. Mandal, Tarit Roy Chowdhury, Gautam Samanta & Dipankar Chakraborti. Analyst, 120, 643-650, 1995.

Arsenic in Ground Water in Six Districts of West Bengal, India: The Biggest Arsenic Calamity in the World. Part 2. Arsenic Concentration in Drinking Water, Hair, Nails, Urine, Skin-scale and Liver Tissue (biopsy) of the Affected People. Dipankar Das, Amit Chatterjee, Badal Mandal, Gautam Samanta, Bhabatosh Chanda & Dipankar Chakraborti. The Analyst, 120, 917-924, 1995.

1996

Arsenic in ground water in six districts of West Bengal, India. Dipankar Das, Gautam Samanta, Badal Kumar Mondal, Chitta R. Chanda, Partha Pratim Chowdhury, Gautam Kumar Basu & Dipankar Chakraborti. *Environmental Geochemistry & Health*, 18(1), 5-15, 1996.

Arsenic in groundwater in seven districts of West Bengal, India-The biggest arsenic calamity in the world. Badal K.Mandal, Tarit Roy Chowdhury, Gautam Samanta, Gautam K.Basu, Partha Pratim Chowdhury, Chitta R. Chanda, Dilip Lodh, Nirmal K.Karan, Ratan K.Dhar, Dipak K.Tamili, Dipankar Das, K.C.Saha & D.Chakraborti. *Current Science*, 70(11), 976-986, June 1996.

1997

Arsenic in groundwater in seven districts of West Bengal, India-The biggest arsenic calamity in the world: the status report up to August 1995. Tarit Roy Chowdhury, Badal Kr. Mandal, Gautam Samanta, Gautam Kr. Basu, Partha P. Chowdhury, Chitta R.Chanda, Nirmal Kr.Karan, Dilip Lodh, Ratan Kr.Dhar, Dipankar Das, K.C.Saha and Dipankar Chakraborti.

Book: *Arsenic: Exposure and health effects*, Edited by C.O.Abernathy, R.L. Calderon and W.R. Chappell, Chapter 9: 91-111, 1997; Publisher: Chapman & Hall, New York.

Non-cancer Effects of Chronic Arsenicosis with Special Reference to Liver Damage. N. Guha Mazumder, J. Das Gupta, A. Santra, A. Pal, A. Ghose, S. Sarkar, N. Cattopadhaya and D. Chakraborti. Book: *Arsenic: Exposure and health effects*, Edited by C.O.Abernathy, R.L. Calderon and W.R.Chappell, Chapter 10:112-123, 1997; Publisher: Chapman & Hall, New York.

Groundwater arsenic calamity in Bangladesh. Ratan K. Dhar, Bhajan Kr. Biswas, Gautam Samanta, Badal Kr.Mandal, D. Chakraborti, Shibtoosh Roy, Abu Jafar, Ariful Islam, Gulshan Ara, Saiful Kabir, A. Wadud Khan, S. Akther Ahmed and S. Abdul Hadi. *Current Science*, 73(1), 48-59, 1997.

Chronic arsenic toxicity in West Bengal. B.K. Mandal, T. Roy Chowdhury, G.Samanta, G.K.Basu, P.P.Chowdhury, C.R.Chanda, D.Lodh, N.K.Karan, R.K.Dhar, D.T.Tamili, D. Das, K.C.Saha and D.Chakraborti. *Current Science*, 72(2), 114-117, 1997.

1998

Impact of safe water for drinking and cooking on five arsenic-affected families for 2 years in West Bengal, India. B.K. Mandal, T.R. Chowdhury, G. Samanta, D.P. Mukherjee, C.R. Chanda, K.C. Saha, D. Chakraborti. *Science of the Total Environment*,

218, 185-201, 1998.

Arsenic levels in drinking water and the prevalence of skin lesions in West Bengal, India. D.N. Guha Mazumder, R. Haque, N. Ghosh, B.K. De, A. Santra, D. Chakraborti and A. H. Smith, *International Journal of Epidemiology*, 27: 871-877, 1998.

Detailed study report of Samta one of the arsenic affected villages of Jessore district, Bangladesh. Bhajan K. Biswas, Ratan K. Dhar, Gautam Samanta, Badal K. Mandal, Imtiaz Faruk, Kazi Saiful Islam, Md. Moniruzzaman Chowdhury, Ashraful Islam and Shibtoosh Roy and Dipankar Chakraborti. *Current Science*, 74(2), 134-145, 1998.

Calcutta's industrial pollution: Groundwater arsenic contamination in a residential area and sufferings of people due to industrial effluent discharge - An eight-year study report. D. Chakraborti, G. Samanta, B.K. Mandal, T. Roy Chowdhury, C.R. Chanda, B.K. Biswas, R.K. Dhar, G.K. Basu and K.C. Saha. *Current Science* 74(4), 346-355, 1998.

1999

Arsenic poisoning in the Ganges delta.

T. Roy Chowdhury, G. K. Basu, B. K. Mandal, B. K. Biswas, U.K. Chowdhury, C. R. Chanda, D. Lodh, S. L. Roy, K. C. Saha, S. Roy, S. Kabir, Q. Quamruzzaman and D. Chakraborti.

Nature, 401, 545-546, 1999.

Groundwater arsenic contamination and sufferings of people in West Bengal, India and Bangladesh: status report up to March 1998. B.K. Mandal, B.K. Biswas, R.K. Dhar, T. Roy Chowdhury, G. Samanta, G.K. Basu, C.R. Chanda, K.C. Saha, S. Kabir, S. Roy and D. Chakraborti.

Book: *Metals and Genetics*, Edited by Bibudhendra Sarkar, Publisher: Kluwer Academic/Plenum Publishers, New York, 3: 41-65, 1999.

Flow Injection Hydride Generation Atomic Absorption Spectrometry for determination of arsenic in water and biological samples from arsenic affected districts of west Bengal, India and Bangladesh. Samanta, T. Roy Chowdhury, B. K. Mandal, B.K. Biswas, U.K. Chowdhury, G. K. Basu, R. Chanda, D. Lodh and D. Chakraborti. *Microchemical Journal* 62, 174-191, 1999.

Arsenic groundwater contamination and sufferings of people in Rajnandgaon district, Madhya Pradesh, India. D. Chakraborti, B.K. Biswas, T. Roy Chowdhury, G.K. Basu, B.K. Mandal, U. K. Chowdhury, S.C. Mukherjee, J.P. Gupta, S.R. Chowdhury, K.C. Rathore. *Current Science*, 77(4), 502-504, 1999.

Arsenic Orphans. Dipankar Chakraborti. Banabithi, Environmental Special Issue, Department of Environment & Forest, Government of West Bengal, India, June, 1999, p7-16.

Possible Arsenic Contamination Free Groundwater Source in Bangladesh. D. Chakraborti, B.K. Biswas, G.K. Basu, U.K. Chowdhury, T. Roy Chowdhury, D. Lodh, C.R. Chanda, B.K. Mandal, G. Samanta, A. K. Chakraborti, M.M. Rahaman, S.Roy, S. Kabir, B.Ahmed, R. Das, M. Salim and Q. Quamruzzaman, J. Surface Sci. Technol. Vol. 15, Nos. 3-4, 180-188, 1999.

Groundwater arsenic contamination and sufferings of people in Bangladesh. U.K. Chowdhury, B. K. Biswas, R. K. Dhar, G. Samanta, B.K. Mandal, T.Roy Chowdhury, D. Chakraborti, S.Kabir, S. Roy.

Book: Arsenic Exposure and Health Effects, Edited by W. R. Chappell, C.O.Abernathy, R.L. Calderon, 165-182, 1999; Publisher: Elsevier, Amsterdam-Lausanne-New York-Oxford-Tokyo.

2000

Groundwater arsenic contamination in Bangladesh and West Bengal-India. U.K. Chowdhury, B. K. Biswas, T. Roy Chowdhury, G. Samanta, B.K. Mandal, G. K. Basu, C.R. Chanda, D. Lodh, K. C. Saha, S.C. Mukherjee, S.Roy, S.Kabir, Q.Quamruzzaman and D. Chakraborti. Environmental Health Perspective, 108: 393-397, 2000.

HPLC-ICP-MS for speciation of arsenic compounds in urine. G. Samanta, U. K.Chowdhury, B.K.Mandal, N.Chandra Sekaran, H. Tokunaga, M. Ando and D. Chakraborti. Microchemical Journal, 65(2): 113-127, 2000.

Arsenic in drinking water and the prevalence of respiratory effects in West Bengal, India. Guha Mazumder DN, Haque R., Ghosh N, De BK, Santra A, Chakraborty D, Smith AH. International Journal of Epidemiology 2000: 29: 1047-1052.

2001

Arsenic groundwater contamination and sufferings of people in West Bengal-India and Bangladesh. U. K. Chowdhury, B. K. Biswas, T. Roy Chowdhury, B. K. Mandal, G. Samanta, G. K. Basu, C.R. Chanda, D. Lodh, K. C. Saha, D. Chakraborti S. C. Mukherjee, S. Roy, S. Kabir, Quamruzzaman.

Book: Trace Elements in Man and Animal, Publisher: Plenum Publishing Corporation, New York, 2001 (in press).

Groundwater arsenic contamination in Bangladesh.

B.K. Biswas, U.K. Chowdhury, M.M. Rahman, K. Paul, G. Samanta, T. Roy Chowdhury, B.K. Mandal, C.R. Chanda, G.K. Basu, D. Lodh, R.K. Dhar, S. Roy, Q. Quamruzzaman, S.C. Mukherjee and D. Chakraborti

Book: Environmental toxicology of metals and metalloids - environmental chemistry, toxicology and health, Australia, 2001 (in press)

Characterization of arsenic bearing sediments in Gangetic delta of West Bengal-India. Dipankar Chakraborti, Gautam K Basu, Bhajan K Biswas, Uttam K Chowdhury, Mohammad Mahmudur Rahman, Kunal Paul, Tarit Roy Chowdhury, Chitta R. Chanda, Dilip Lodh

Book: Arsenic Exposure and Health Effects, Edited by W. R. Chappell, C.O.Abernathy, R.L. Calderon, 2001; Publisher: Elsevier, Amsterdam-Lausanne-New York-Oxford-Tokyo.

Groundwater arsenic calamity in West Bengal-India and Bangladesh. U.K. Chowdhury, B.K. Biswas, G. Samanta, B. K. Mandal, T. Roy Chowdhury, R. K. Dhar, G. K. Basu, C. R. Chanda, K. C. Saha, S. Roy, S. Kabir and D. Chakraborti.


Book: Bioavailability and its potential role in risk assessment, Publisher: Oxford, 2001 (in press).




Skin lesions or symptoms observed in Bangladesh and West Bengal (taken by Dr. Chakraborti)



Arsenic affected districts of Bangladesh



Endemic states for fluorosis of India



note)

The above 3 figures are showing the distributions of affected populations with arsenic or fluoride in West Bengal, Bangladesh and India, respectively. It is noteworthy that there are areas affected with both of them especially in West Bengal and Bangladesh.

1.2 Chronic arsenicism in Ronphibun, Thailand

~ past, present and future

Shoko Oshikawa¹, Alan Geater¹, Virasakdi Chongsuvivatwong¹,

Tada Piampongsan²,

Dipankar Chakraborti³, Gautam Samanta³, Badal Mandel³,

Nobuyuki Hotta⁴, Yasuko Kojo⁴, Hiromi Hironaka⁴

1. Epidemiology Unit, Faculty of Medicine, Prince of Songkla University, Thailand

2. Institute of Dermatology, Bangkok, Thailand

3. School of Environmental Studies, Jadavpur University, India

4. Asia Arsenic Network, Japan

PAST


Ronphibun



Ronphibun Sub-district,
Ronphibun District,
Nakhon Si Thammarat
Province, Thailand

Source of map:

JICA report in 2000



Affected area

- Historical tin mining area
- Waste from mining activities deposited throughout the area.
- Environmental contamination by arsenic had been found in surface - and under ground - water, and soil.

(photograph1~ previous dredging site)

Affected people

- “Kai-dam” was endemic but cause was generally unknown locally.
- Most of residents had used shallow well water.
- In 1987, kaidam was officially recognized as chronic arsenicism when one woman was diagnosed at the Institute of Dermatology in Bangkok.

Governmental survey in 1987

- Ministry of public health set a station survey for clinical examination at Ronphibun District Hospital.
- Subsequently 1151 cases were diagnosed as chronic arsenicism and the severity of their lesions was classified into stage 1 to 4.*¹
- The cases in severe stage were treated.
- Water in a large number of shallow wells was found to have high arsenic content (>0.05~4.45 mg/l), and the distribution of contaminated wells scattered not-homogeneously in the area.*²


*¹ The Report on Rectification of Arsenic Poisoning Problem in Ronphibun, Ronphibun District, Nakorn

si thamarat Province: Nakorn si Thamarat Provincial Health Office, 1988.

*² The Office of National Environmental Board. *Report of Arsenic Contents of Well Waters in Ronphibun*

District, Nakorn si thamarat Province, Bangkok: National Environmental Board, 1987.

Prevalence of shallow wells contaminated with arsenic (>0.05mg/l) by village in 1987)



The data from: Rodklai A, Shindell S. *Chronic arsenic poisoning in Nakorn si thammarat*. Ninth Report in Field Epidemiology Training Project, Bangkok. Undated.

Intervention programmes, 1987 ~

- Distribution of large water jars for storage of rain water
- Installing deep well
- Improvement and expansion of piped water supplies
- Closure of shallow wells more heavily contaminated with arsenic

*Prevalence survey in 1994 *¹*

- Prevalence of subjects with chronic arsenicism was 26.3% in Ronphibun Sub-district.
- Prevalence varied among the villages.
- Higher prevalence was found in village 1 (53%), village 12 (50%) and village 5 (36%).


- 17.6% of households were still using shallow well water for drinking due to no supply of piped water.

*¹ .Rodklai A. *Prevalence Survey of Chronic Arsenic Poisoning and Health Status of the People in Ronphibun, 1994*. Report of an Epidemiological Survey in Ronphibun, Thailand, 1994.

Here are 2 figures showing the prevalence.



1. Prevalence of As contaminated wells (>0.05 mg/l) by village (1987)

*Data from ; Rodklai A, Shindell S. *Chronic arsenic poisoning in Nakorn si thammarat*. Ninth Report in Field Epidemiology Training Project, Bangkok. Undated.



2. Prevalence of subjects with skin lesions by village in 1994

. *Data from ; Rodklai A. *Prevalence Survey of Chronic Arsenic Poisoning and Health Status of the People in Ronphibun, 1994*. Report of an Epidemiological Survey in Ronphibun, Thailand, 1994.



CURRENT SITUATION : STUDIES IN 1997

Study 1~ changes in skin lesions severity following intervention in 1987

- Eight hundred and eighteen subjects with arsenical skin lesions first examined in 1987 and having address in Ronphibun sub-district were followed-up.

Study objectives:

- To describe changes in stage of skin lesions over the 10-year period since 1987
- To explore the role of drinking water in the changes

Methodology

- Questionnaire-based interview (sociodemography, history of residence, treatment, mining, water use)
- Visual assessment and photographic record of skin lesions
- Arsenic analysis of drinking water in the field
- Observation of well and water supply facilities
- Staging of skin lesions

Staging

The staging classification used in 1987 survey, was one proposed by the Institute of Dermatology, Bangkok.

Stage 0: no lesions (but may have elevated hair, nail, urine and/or blood As levels)

Stage 1:*palms/soles* Small pigmented macules or a few pin-headed dermal papules

Stage 2:*hands/feet* Multiple pin-headed dermal papules and/or large keratotic papules (>5mm diameter)

Stage 3:*hands/feet* Stage 2 lesions with pre-Bowen;s disease

Stage 4: *any site* Bowen's disease, basal cell carcinoma, squamous cell carcinoma

The results of study


Follow-up status

- Out of 818 subjects, 334 (40%) could be followed-up and the remainder had died (35), migrated (193), were temporary absent (55) or were unknown (201).


Change in stage over the 10 years

- Both regression and progression of skin lesions had occurred.
- Minimum proportions undergoing regression, assuming none of the missing had regressed, would be 29%.
- Minimum showing progression, assuming none of the missing had progressed, would be 5%.
- Among the 4 subjects who had progressed from stage 2 to stage 4, one had developed basal cell carcinoma and three Bowen's disease in 1997.
- One of 2 subjects who had Bowen's disease in 1987, had developed squamous cell carcinoma.


Change in drinking water: 1986 and 1997



Change in cooking water : 1986 and 1997



Change in bathing water : 1986 and 1997



Analytical methodology

- Used BINARY LOGISTIC REGRESSION to explore the relationship between pattern of drinking water and changes of skin lesions.
- Stratified on age and village of residence.
- Confined to stages 1 and 2 subjects in 1987, using separate models.
- The model was adjusted for occupation in mining.

The role of drinking water in the changes

- The use of rain water for drinking compared that of piped or bottled water has not been fully effective in reducing the severity of skin lesions.

Study2 ~ lesion type and severity

Among the followed-up subjects, a detailed description of cutaneous symptoms of chronic arsenicism was made in 203 subjects. The skin lesions showed;

- Wide variation in skin lesions was apparent.
 - Type ~ hyperkeratosis, hyperpigmentation, depigmentation
 - Distribution ~ spotty, diffuse
 - Site ~ trunk, limbs, palms, soles, etc.
 - Severity

Study objectives

- To describe skin lesion profiles.
- To develop a preliminary model to explore relationship between exposure mode and lesion profile.

Methodology ~ detailed description of skin lesions

Physical examination done by a dermatologist specialized in these lesions.

- whole body observation ~ 16 sites
- using standardized form
- pigmentation, depigmentation and keratosis
- 4 levels of severity (-, +, ++, +++)
- Bowen's disease and skin cancer presence based on histopathological examination

Results

- Spotty pigmentation, diffuse pigmentation, spotty keratosis, diffuse keratosis and depigmentation were observed in 84%, 54%, 74%, 55% and 42%, respectively.
- Bowen's disease, basal cell carcinoma and squamous cell carcinoma were diagnosed in 12, 2 and 1 subjects, respectively.
- Preliminary analysis to explore relationship between exposure mode and lesion profile is going on.

FUTURE ~ WHAT STILL NEEDS TO BE DONE?

- Further studies on several aspects of arsenic contamination in Ronphibun are still needed.
- How can countermeasures be improved to reduce exposure to arsenic?

Further studies

- Continued follow-up of the subjects with chronic arsenicism.
- New case finding.
- Clarification of the role of drinking water.
- Exploration of possible pathways of contamination other than through drinking contaminated water.

Countermeasures

- Completion of the safe water supply in the contaminated area.
- Empowerment of residents to participate in decision-making regarding pollution abatement.
- Declaration of the contaminated area as a “Pollution Controlled Area” in order to officially accord high priority to pollution control measures.

1.3 Comments and Recommendation

An arsenism study on communities in Bangladesh

Chiho Watanabe

School of International Health, University of Tokyo

< See the slides on the next pages for the contents. >

Background

Bangladesh

population: 140,000,000 (700-800/km²)
tubewell: >95% tubewell for drinking water
Risk population in BD: 35,000,000
first report of As patient: 1994

1

Comments and Recommendations

An Arsenism Study on Communities in Bangladesh

Chiho Watanabe and Ryutaro Ohtsuka
School of International Health
University of Tokyo


2

Focus

- * Mazumder et al. (1998) West Bengal n=ca. 7,700
skin lesions males>females
malnutrition enhanced the symptom
- * Tondel et al. (1999) n=ca. 1,500
skin lesions males>females
- * Ahsan et al. (2000) (n=167) males<>females [As]u

Dose-response relationship using biological media
Identification of modifying factors/sensitive population
* Tseng (1977) Taiwan
male > females (skin cancer) males <> females

3



Two small communities, SV and SP, in northwestern Bangladesh were the subjects of this study.

4



Sadasibpur, Nawabganj Fall 1998

5



Serpur Vander, Nawabganj (spring 1999)

6




*Flooded ricefield Serpur Vander, August 1998
Almost the same area shown in the previous picture*

7



8



*Although some clustering can be seen, the geographical distribution
As concentrations of the tubewells showed a patchy pattern, and
appeared hardly predictive.*

9

Table 2. Age, nutritional status, and As exposure in the groups divided by sex and by area *


area	sex	Age ^a [yr]	Height [cm]	Body Weight [kg]	Fat ^b Weight percent [%]	[As] _{ur} ^a [mg/L]	[As] _{ur} ^a [mg/gCr]
SV	male	36±1.5 (112)	162±5 (111)	52±7 (111)	10.7±1.4 (111)	28±6.0 (104)	204±2.1 (64)
SV	female	36±1.4 (138)	151±6 (137)	43±6 (138)	21.4±1.2 (137)	26±6.6 (126)	219±2.4 (108)
SP	male	40±1.5 (108)	164±6 (107)	51±7 (100)	10.7±1.5 (108)	11±6.6 (108)	126±2.3 (69)
SP	female	35±1.5 (193)	152±5 (192)	44±7 (162)	20.9±1.3 (192)	10±6.4 (189)	174±2.5 (121)

ANOVA--


	Area	ns	#	ns	ns	***	***
'Sex	#	***	***	***	***	ns	#
Area*Sex	#	ns	ns	ns	ns	ns	ns

- NOTES:
- Geometric mean and SD are shown.
 - Calculated from skinfold thickness.
 - Two-way ANOVA using log-transformed variables.
 - *** $p<0.001$, # $0.05<p<0.1$, ns = not significant ($p>0.1$).


10



11



12



1 3

Table 3. Stepwise multiple regression analyses of urinary arsenic excretion ([As]u)^{*1}

area	sex	adjusted -R ²	intercept	c	[As] tw	P	Age	fat
SV	mal	0.47	3.486	***	0.294	***	-0.293	+ ^{*2}
	fem	0.43	2.839	***	0.320	***		
SP	mal	0.64	3.515	***	0.356	***		
	fem	0.51	3.482	***	0.328	***		-0.440 *

NOTES:

*1; Parameter estimate and its significance are shown.
 *2; A blank means that the parameter was not selected (entered) as a significant independent variable in the regression model.
 *** p<0.001, * p<0.05, + 0.05<p<0.1.

1 4

Table 1. Classification criteria of the dermatological stage (DS) and distribution of the subjects by DS

Criteria for the stage ^a		Frequency distribution ^b			
DS	Rank	Manifestation	Prevalence [%] ^c (total = 468)	n	% of total
0	(none)			213	45.5
1	1	sole 1	53.8	148	31.6
2	2	sole 2	18.4	37	7.9
3	3	palm 1	9.8	43	9.2
	4	trunk 1	8.3		
4	5	sole 3	5.1	19	4.1
	6	palm 2	3.4		
5	7.5	sole 4	1.1	8	1.7
	7.5	trunk 2	1.1		
	9	palm 3	0.6		

For an individual diagnosed as sole = 3, palm = 2, trunk = 1, the corresponding ranks are 5, 6, and 4, respectively. The DS corresponding to the highest of these three ranks (=6), which is 4, is assigned as this individual's DS (see text for detail).

b. Actual distribution of the subjects classified for each DS.
 c. Prevalence of the manifestation


1 5

% Distribution of the Dermatological scores (Feb 199)

Grade	total	By village		By gender (p<0.001)	
		SP	SV	Males	Females
0	55	55	54	44	62
1	23	26	18	24	23
2	13	12	15	17	11
3	5	4	8	8	3
4	3	2	3	6	0
5	1	1	1	1	1
total	100	100	100	100	100
(N)	561	353	208	238	323

(Including those less than 20 yrs old)

1 6



1 7

Urinary arsenic
 good correlation with twAs
 stable
 reflect recent exposure

Dose-response relationship
 twAs determined uAs; little effect of sex, age, fat
 females more resistant to chronic As

Other modifying factors
 nutrition, micronutrients as modifying factors
 effects on development

1 8

Session 2

Current Status of Arsenic Pollution and Health Impacts in China

2.1 The Current Situation of Endemic Arsenism in China

Guifan Sun

School of Public Health, China Medical University, P.R.China

Since the endemic arsenism was first found in the mainland of China in 1983 in Xinjiang Province, there have been more cases identified in Inner Mongolia, Shanxi and Guizhou Province. Total population exposed to arsenic has been over two million and the diagnosed patients came to 20 thousand. Now the epidemic area is still expanding. The etiology of arsenism in China is mainly divided into two types: 1. Drinking water type: Many people drink water from pump well 20~ 30m deep underground rich in arsenic from 0.2mg/L to 2.0mg/L, maximum up to 4.44mg/L. 2. Burning coal type: People use the coal containing high concentration arsenic as fuel for cooking, keeping warm and drying grains and peppers in kitchen, so the food and indoor air are seriously polluted by arsenic. The average arsenic concentration in coal is 876.3mg/kg. In polluted kitchen air is 445ug/m³. In corns and peppers are 4.13 and 512.0mg/kg respectively. Although some preventive measures have been taken, how to remove arsenic from drinking water and prevent burning coal type arsenism is still an important problem now in China.

Distribution

In China, chronic endemic arsenism was first found in Jiayi and Tainan, Taiwan province in 1968. The area of the disease was about 300km² and the population was about 150 thousand. It is well known as the “Black Foot” disease.

However, the disease in the mainland of China was reported only in recent years. In the mainland, the disease was first found in Kuitun, Xinjiang province in 1983. The patients were more than 2000 and the population of the epidemic area was about 100 thousand.

In 1989, a large-scale survey was conducted in Chifeng, Bayanzhuoermeng district and Huhhot of Inner Mongolia Autonomous Region. The diagnosed patients were 1774, the population involved reached 600 thousand.

It was not until 1994 that arsenism was found in Datong Basin, Shanxi province. The epidemic area was over 900 km² and the exposed population were up to one million. Arsenism in all the places mentioned above was induced by water rich in arsenic.

In 1991, 2600 patients and 200 thousand population were found once again in six

cities or counties of Guizhou province. The use of high arsenic coal was the reason of arsenism in this area.

By the end of 1997, it is estimated that the total population exposed has been over two million and the diagnosed patients has come to over 20 thousand. What's more , the epidemic area is still expanding.

Although the history of arsenism in the mainland is short, the situation and the spread of the disease are very serious.

Chronic endemic arsenism in China can be divided into two types according to the exposure sources. The first type is caused by drinking water, in which arsenic concentration is very high. The second type is caused by burning coal.

Etiology

1. Drinking water type: In the 1970's, with the improvement of peasants' living conditions, the surface water is substituted by the pump well water to improve the water quality and to prevent the endemic fluorosis. Many people drink water from pump well 20- 30m deep under the ground, which is unexpectedly rich in arsenic from 0.2mg/L to 2.0mg/L, maximum up to 4.44mg/L. It is reported that in some areas, the average arsenic concentration of drinking water is tens of times higher than the 0.05mg/L China standard.

2. Burning coal type: The local inhabitants use coal containing high arsenic as fuel for cooking , keeping warm and drying grains and peppers in kitchen, so that the food and the indoor air are seriously polluted by arsenic. It is a unique exposure of China , where the average arsenic concentration in coal is 876.3mg/kg, maximum concentration up to 9600mg/kg. The average arsenic concentration in kitchen air is $445\mu\text{g}/\text{m}^3$, which is over 150 times higher than the hygienic standard of China. The arsenic concentration of the indoor floating dust is 3800mg/kg, which is 57 times as much as that of the control. The arsenic concentration of corns and peppers are 4.13 and 512.0mg/kg respectively. Recently, as the coal consumption grows continuously, the patients of burning coal type have been increasing sharply.

In the initial period of hyperkeratosis, we can see the hidden papules under the skin of the palms and the soles. The papules are semi-transparent in early phase, then keratinize to be helomaform, verruciform, rhagadiform and cauliflower-likeform. In the seriously affected patients, the hyperkeratotic skin fuse together, appearing to be withered tree roots or toad-like skin.

Chronic arsenic exposure can also induce the Bowen's disease. It mainly happens in long-term exposure areas. It can be solitary or multiple and may be found before the

cancer or simultaneously with it. There is no difference in pathology between arsenic skin cancer and other skin cancers. The situation has happened in Huhhot of Inner Mongolia.

In addition to these peculiar signs, other arsenic-induced symptoms are all prevalent in the epidemic areas, such as fatigue, palpitation, numbness of hands and feet, abnormal sensation, skin capillarioscopy and peripheral circulation disturbance.

It is common for more than one members of a family being affected.

Preventive measure

Arsenism not only influence people's health, work and family happiness, but also bring them unimaginable pains in both soul and heart. The government has been paying high attention to endemic arsenism. In July 1992, arsenism was officially regarded as one of the eight endemic diseases which are administrated by the National Endemic Disease Office. A lot of medical groups and researchers were sent, special-purpose funds was allocated by the government to prevent and control the disease. The underground water has already been substituted by the tap water in Xinjiang. A large-scale water improvement project has been carried out in Inner Mongolia and Shanxi province. In Guizhou, the local government control the exploitation and use of high arsenic coal strictly, publicize the harmfulness of arsenic coal, and help the local residents rebuild the stoves. But there are a lot of work need to be done for prevention and treatment.

Up to now, only some villages have altered water supported by government and most of them are still use the polluted drinking water. Although some equipments of removal of arsenic were tested in some places, most of them have been stopped in use because they were both inconvenient and expensive. How to remove arsenic from water is still to be solved.

In order to organize the country's investigation and for deep researches, and to make cooperation with international organization and foreign experts, the Fluoride and Arsenic Society of China (FASC) was founded in 1996. In October 1998, the specialists all over the country gathered by the Fluoride and Arsenic Society of China (FASC) to discuss and make clinical diagnostic standard of arsenism. And in August 1999, Pan Asia-Pacific Conference on Fluoride and Arsenic Research was held in China. Now FASC is just making the preparation for the general survey throughout the country.

2.2 Progress on Arsenism Research in China

Jin Yinlong

IEHE, CAPM, China

< See the slides on the next pages for the contents. >

Progress on Arsenism Research in China

Prof. Jin Yinlong
Director
Institute of Environmental Health & Engineering, CAPM

Slide 1 of 31

Institute of Environmental Health and Engineering, CAPM

1

- Introduction
- Current state of arsenism research
- Unsolved problems
- Future studies

Slide 2 of 31

Institute of Environmental Health and Engineering, CAPM

2

1. Introduction

Arsenic is widely distributed in the environment and chronic consumption of arsenic-contaminated drinking water or coal is epidemiologically linked to many toxic effects. There are two types of endemic arsenism. In China, one is called drinking water type, the other is named as coal burning type.

Slide 3 of 31

Institute of Environmental Health and Engineering, CAPM

3

Our government has also paid much attention to arsenism. It was taken as one of the main preventive endemic diseases by Ministry of Health in 1992, and its preventive program was also listed in the Ninth Five-Year Plan in 1996.

Slide 4 of 31

Institute of Environmental Health and Engineering, CAPM

4

2. Current state of arsenism research


A great-scale investigation on arsenism has been conducted since 1980's, the situation, characteristic and principia of arsenism are almost clear now. The mechanism of arsenism also has been studied, for example, arsenic induces peroxidation, affects on the activities of protein, enzyme and nucleic acid, damages biomembrane and cell. On the other hand, many interfered studies by changing drinking water or improving stove have been carried out in arsenism areas.

Slide 5 of 31

Institute of Environmental Health and Engineering, CAPM

5

1) Distribution of arsenism in China



Slide 6 of 31

Institute of Environmental Health and Engineering, CAPM

6

2) Situation of arsenism areas (1998)

Type	Area	Concentration of arsenic (mg/L)	Arsenism patients (case)	Exposed population (10,000)
	Xinjiang	0.05-0.85	2000	10
Drinking water	Inner Mongolia	0.129-0.596	2455	22
	Shanxi	0.03-1.41	3000	100
Coal burning	Guizhou	26.5-2116.7 mg/Kg	2600	20
Total			10,055	192

Slide 7 of 31

Institute of Environmental Health and Engineering, CAPM

7

3) Characteristic of arsenism disease

Hyperpigmentation, hypopigmentation and keratosis are the most important characteristics of arsenism disease in China mainland, but it is blackfoot disease in Taiwan. There are some disputes on the pathological changes:

Slide 8 of 31

Institute of Environmental Health and Engineering, CAPM

8

- Some authors suggested that the clinic behaviors of arsenism started with keratosis in palm like papilla, then progressed to hyperpigmentation and hypopigmentation.
- Some authors suggested that the disease started with hyperpigmentation and hypopigmentation clinically, it progressed to keratosis only when people drink water with high enough concentration of arsenic

Slide 9 of 31

Institute of Environmental Health and Engineering, CAPM

9

4) Dose-response relationship


Arsenism is mostly occurred in the people who drinks water or combusts coals with high content of arsenic. It has as obvious family assemble and region assemble characteristic.

Slide 11 of 31

Institute of Environmental Health and Engineering, CAPM

10

Relationship between concentration of arsenic in drinking water and incidence rate of arsenism in Xinjiang




Slide 12 of 31

Institute of Environmental Health and Engineering, CAPM

11

Relationship between concentration of arsenic in drinking water and incidence rate of arsenism in Inner Mongolia




Slide 13 of 31

Institute of Environmental Health and Engineering, CAPM

12


Relationship between concentration of arsenic in coal and incidence rate of arsenism in Guizhou



Slide 14 of 31 Institute of Environmental Health and Engineering, CAPM

1 3

Comparison of incidence rate of arsenism in different age groups in Xinjiang (concentration of As is 0.75mg/L, drink for 5 years)



Slide 15 of 31 Institute of Environmental Health and Engineering, CAPM

1 4

5) Related to other diseases

Region	Cardiovascular		Cancer		
	Abnormal cardiograph (%)	Renal syndrome (%)	Digestive (%)	Respiratory (%)	Skin (%)
Xinjiang	78.9	34.7	17.4	52.2	
Inner Mongolia	80.0	69.0	35.9	38.5	14.1

Slide 16 of 31 Institute of Environmental Health and Engineering, CAPM

1 5

6) Mechanism research

Study on the relation between peroxidation and arsenism in Inner Mongolia


The results show :

Arsenic in body can induce lipoprotein peroxidation and produce free radical as well as decrease the activity of SOD, GSP-Px. As a result damage people's health.

Slide 17 of 31 Institute of Environmental Health and Engineering, CAPM

1 6

Comparison of SOD and MDA in blood in arsenism case and control group in Inner Mongolia



Slide 18 of 31 Institute of Environmental Health and Engineering, CAPM

1 7

Study on interaction between arsenic and human erythrocyte by FT-IR


The results show :

Arsenic can affect the conformation and configuration of membrane phospholipid, decrease its fluidity, and stimulated peroxidation, as a result, damage the erythrocyte. On the other hand, arsenic can also change the secondary structure of membrane protein, affect its stability and function.

Slide 19 of 31 Institute of Environmental Health and Engineering, CAPM

1 8

FT-IR Spectra of membrane phospholipid of human erythrocyte (1)




Slide 20 of 31

Institute of Environmental Health and Engineering, CAPM

1 9

FT-IR Spectra of membrane phospholipid of human erythrocyte (2)




Slide 21 of 31

Institute of Environmental Health and Engineering, CAPM

2 0

Effect of As(III) on secondary structure of membrane protein



Slide 22 of 31

Institute of Environmental Health and Engineering, CAPM

2 1

7) Prevention and therapy of arsenism

Therapy by drugs :

- 1, **DMSA**: increase the excretion of excessive arsenic in body as well as improve microcirculation
- 2, **Se**: increase the activity of GSH-P_x and Na⁺-K⁺-ATPase and improve microcirculation.

Prevention :

- 1, Search water with low concentration of As, eliminate excessive arsenic in body in water.
- 2, improve the stove and decrease the concentration of As in indoor air.
- 3, Study on the physical and chemical method for elimination of arsenic.

Slide 23 of 31

Institute of Environmental Health and Engineering, CAPM

2 2

3. Unsolved problems

1. Various criterion :

There were many studies of arsenism in China before, but they base on different diagnostic criterion, and different determination method for arsenic. So some results in different regions are not comparable.

Slide 24 of 31

Institute of Environmental Health and Engineering, CAPM

2 3

2. The distribution of arsenism is unclear :

Major researches of arsenism are mainly carried out in north three provinces, such as Xinjiang, Inner Mongolia and Shanxi province. While researches in south of China just have been began, for example in Huan and Jiangxi province. According to the National Survey on Drinking Water Quality and Water-born Diseases, there were total 5,630,000 people, who drink high arsenic content water (0.05mg/L). It seems that arsenism distributes in other regions beside Xinjiang, Inner Mongolia, Shanxi and Guizhou.

Slide 25 of 31

Institute of Environmental Health and Engineering, CAPM

2 4

3. Limited research on mechanism of arsenism :

The mechanism of arsenic toxicology is unclear, especially the bi-effects and carcinogen of arsenic. Many studies suggested that arsenism is associated to skin cancer, lung cancer, bladder cancer, and so on. However, it is necessary to study further in the molecular level.

Slide 26 of 31

Institute of Environmental Health and Engineering, CAPM

2 5

4. Future studies

Arsenism is not only located in Xinjiang, Inner Mongolia, Shanxi and Guizhou province in China, but also in other regions, such as Ningxia, Shaanxi, Beijing, Jilin, Liaoning, Hebei and Henan province for drinking water type arsenism, Xichuan, Yunnan, Guangxi, Hunan, Hubei and Jiangxi province for coal burning type arsenism. To see the distribution and harm of arsenism in china, our government decides to carry out a national survey on arsenism.

Slide 27 of 31

Institute of Environmental Health and Engineering, CAPM

2 6

Study protocol

- 1、 Organize a team of national investigation on arsenism
- 2、 Reported the concentration of As in water and coal and situation of arsenism in each region by the charge department of endemic disease.
- 3、 According to the reported information, select the typical regions of arsenism.
- 4、 Unify design and survey base on the standard for determining arsenism region caused by arsenic in environment and standard for diagnosing arsenism.

Slide 28 of 31

Institute of Environmental Health and Engineering, CAPM

2 7

Content of studies

Investigation of environmental factor :

determine the concentration of As in water , soil, food, coal, indoor air and some biologic materials in selected regions

Investigation of arsenism :

investigated the population data, situation of drinking water, situation of coal consumption, residence condition, history of disease and life style base on the standard for determining arsenism region caused by arsenic in environment and standard for diagnosing arsenism.

Slide 29 of 31

Institute of Environmental Health and Engineering, CAPM

2 8

Purpose

- 1、 Map the distribution of arsenic in drinking water and coal in whole country.
- 2、 Map the distribution of arsenism areas, exposed population and arsenism patient in whole county.

Slide 30 of 31

Institute of Environmental Health and Engineering, CAPM

2 9

2.3 Assessment of Arsenic Removal from Drinking Water by New Adsorbents

He Gongli, Yang Jiaolan, Yu Xiaoyi and Yue Yingling
(Institute of Environmental Health and Engineering, CAPM,
Nan Wei Road, Beijing, 100050,China)

SUMMARY: Efficiency comparison of arsenic removal from drinking water by cerium hydroxide, magnum oxide, calcium oxide and Cu-impregnated activated carbon has been carried out. Effect of contact time on the efficiency of arsenic removal by cerium hydroxide and Cu-impregnated activated carbon has been investigated. The removal of arsenate ion by coprecipitation with cerium hydroxide and Cu-impregnated activated carbon for 24 hr was more than 98% at neutral pH. The contact time effects evidently the efficiency of arsenate ion removal.

KEYWORDS: *arsenate, coprecipitation, cerium hydroxide, and Cu-impregnated activated carbon*

Introduction

Arsenic is a toxicant with widespread biologic effects. It has been defined as priority consideration carcinogen by the Center of Disease Control of the United States (CDC) and International Association for Research Cancer (IARC) ^[1]. The hygienic standard of arsenic in drinking water is turning the stricter on International. Japan and German reduced the standard level of arsenic in drinking water from 0.05mg/L to 0.01mg/L in 1993 and 1996, respectively ^[2-3]. Pontius^[4] reported that a new rule for arsenic in drinking water was announced in the United States in 1992. Preliminary calculation for the maximum contaminant level (MCL) suggest that a value of 0.002mg/L will likely be proposed, which is a 25-fold decrease from the present interim MCL value of 0.05mg/L. If this new MCL is adopted, arsenic could become a new substance of concern and it may be a challenging task for treatment technology in the future.

Major treatment methods to remove oxyanions of arsenic include complexation with polyvalent metal species such as ferric iron and coprecipitation with amorphous hydrous metal oxides, lime softening, ion-exchange and activated alumina. The conventional methods of arsenic removal such as coagulation precipitation with lime, alum, or ferric sulfate produce a wet bulky sludge and often require final filtration for secondary treatment. Recent literature shows that various hydrous oxides of rare earth elements exhibit high-adsorption capacity for anions. Various adsorption studies of metal ions have been done on silica-water interface. The materials improved with rare

earth elements can remove 99.9% of arsenate from aqueous solution [5]. Which prompted that some typical mechanisms of interact can exit between rare earth elements and arsenate.

The concept of multiple separation by chemisorptive filters was applied and investigated in the process of arsenic removal from water. Chemisorption filters were made by the paper manufacturing method and consisted of cellulose, cationic and anionic ion exchangers, activated carbon and a corresponding chemical agent. Chemisorption filters were activated with Ag^+ , Mg^{2+} , Cu^{2+} , Al^{3+} and Fe^{3+} ions. The best result in arsenic bonding was obtained when Cu^{2+} ions were applied for chemical activation of activated carbon. The effectiveness of this adsorbent was 0.18mmol/g [6].

In the present study, metal oxides were used for the removal of arsenate ion by coagulation precipitation in aqueous solution at different pH and contact time. The kinetic study of coagulation precipitation of arsenic was conducted. The objectives of this work are (1) to evaluate the efficiency of arsenic removal from drinking water by different adsorbents and (2) to determine the optimum operation parameters affecting the removal of arsenic by cerium hydroxide and Cu-impregnated activated carbon, such as raw water pH values, contact time and initial concentrations of arsenate.

Materials and Methods

Chemicals and materials used in this study included:

Ce(OH)₄: cerium hydroxide (prepared by our laboratory)

MgO: magnum oxide (analytical grade)

CaO: calcium oxide (analytical grade)

Cu-AC: Cu-impregnated activated carbon (prepared by our laboratory)

Arsenic solution with different concentrations was prepared by dissolved dibasic sodium arsenate in deionized water.

Cerium hydroxide preparation: dissolved 5g Ce(SO₄)₂ into 100ml deionized water, add 2%NaOH solution(excessive), producing a yellow precipitate that was filtered with filter paper, washed with deionized water to neutrality, and dried it in drier. A yellow powder product was obtained.

Cu-impregnated activated carbon preparation: To 100ml of 5% CuSO₄ solution was added 40g of powder activated carbon (40 mesh). The mixture was shaken for 24 hours at room temperature and filtered with filter paper. The residue was washed repeatedly with deionized water and dried it at room temperature.

Coagulation precipitation of arsenic in a batch experiment: 30mg of four adsorbents (see Table 1) were suspended, respectively, in 100ml of arsenic-containing

water with continuous stirring. After mixing for 1-2 min, followed by static reaction at room temperature for 24 hours. The supernatant was drawn off by suction for determination of pH and arsenic. The arsenic concentration in the raw water and treated water was determined by GFASS method.

Results and Discussion

The main findings of the different adsorbents examined are given in the Table 1. Four adsorbents are effective for As^{5+} removal. But a disadvantage of MgO and CaO is that the pH value in the treated water is about 10-12. It is too high as drinking water use.


Effect of contact time on the efficiency of arsenic removal by cerium hydroxide and Cu-impregnated activated carbon: Removal of arsenate ion at different contact time at raw water pH 7.16 is summarized in Figure 1. The percentage removal of arsenate ion increased as the contact time increased. The removal was found to be more than 98% at 24 hr contact time. When contact time decrease from 24hr to 4 hr, removal of arsenate ion decreased from 99.0% to 60.4% for cerium hydroxide and from 98.7% to 61.3% for Cu-impregnated activated carbon.

Effect of initial concentrations of arsenic on the efficiency of arsenic removal by cerium hydroxide and Cu-impregnated activated carbon: Removal of arsenate ion by cerium hydroxide and Cu-impregnated activated carbon at different pH and initial concentrations are summarized Figure 2a and 2b. The removal was found to be more than 99% for both adsorbents of cerium hydroxide and Cu-impregnated activated carbon in the final pH range of 4.5 to 7.5 at the initial concentration of 0.52mg/L. The percentage removal of arsenate ion decreased as the initial concentration of arsenate ion increased. Removal of arsenate ion decreased from 99.6% to % for cerium hydroxide, and from 99.6% to % for Cu-impregnated activated carbon with an increase in concentration from 0.52 to 1.98mg/L. Arsenate can be removed down to 0.002mg/L when the initial concentration of 0.52mg/L was taken.

Table 1. Efficiency of As⁵⁺ removal of different adsorbents
For 100ml of water over 24 hr at room temperature

Group	Adsorbents and amounts (mg)	Raw Water		Treated Water		
		As(mg/L)	pH	As(mg/L)	Removal(%)	pH
A	Ce(OH) ₄ -30	0.514	4.49	0.002	99.6	6.17
		0.531	7.16	0.005	99.0	7.20
		0.526	9.18	0.022	95.8	7.94
B	MgO-30	0.514	4.49	0.022	95.7	9.89
		0.531	7.16	0.036	93.2	10.5
		0.526	9.18	0.045	91.4	10.7
C	CaO-30	0.514	4.49	0.021	95.9	11.5
		0.531	7.16	0.031	94.2	12.6
		0.526	9.18	0.039	92.6	12.8
D	Cu-AC-30	0.514	4.49	0.002	99.6	5.32
		0.531	7.16	0.007	98.7	6.89
		0.526	9.18	0.017	96.8	8.45

Figure 1. Effect of contact time on the efficiency of arsenic removal by Ce(OH)₄ and Cu-AC






Figure 2 Removal of arsenate ion by cerium hydroxide and Cu-impregnated activated carbon as a function of pH and initial concentration. Adsorbents amount was 30mg; contact time, 24hr; and temperature, 20 .
2a cerium hydroxide; 2b Cu-impregnated activated carbon

Reference

- [1] Jeffer A L., Robert FS, Christopher TD, environmental toxicants. Priority Health Conditions, 1993, 77
- [2] 齊藤隆彦, 浄水処理技術中逆向連続流動床式過濾飲水除 方法, 公開特許公報 (A), 1995 年, 特開平 7-88482
- [3] Hoerner G., Remove of arsenic experiences of commercial plants, DVGW-Schriften,Wasser, 1993, 82: 189
- [4] Pontius FW, A current lood at the federal drinking water regulations, J. Am Water Works Assoc., 1992,84: 36
- [5] Syed AW., Haron J., Akira V., et al. Removal of arsenite and arsenate ions from aqueous solution by basic yttrium carbonate, Water Research, 1996, 30(5): 1143
- [6] Ljubinka VR, Milan MM., Arsenic removal from water by chemisorption filters, Environmental Pollution, 1992, 75: 279

2.4 Comments and Discussion

A Study of arsenism via consumption of arsenic contaminated ground water and intervention study in water mitigation in Inner Mongolia.

Takahiko Yoshida

Asahikawa Medical College

To study a relation between arsenic (As) exposure and As related skin lesions, field researches were done on villagers who drunk naturally As contaminated well water in Inner Mongolia. Then to evaluate a reversibility of skin lesions associated to As exposure, interventional study by providing drinking water with lower concentration of As in mitigation were conducted as periodical follow up studies after changing water source. Correlation between As exposure index or As exposure duration and skin lesions were analyzed on individual villagers. Skin symptoms were obvious in male than female, especially in skin dyspigmentation. As exposure index were related with hyperkeratosis on palm in both adult and children. While hyperkeratosis on sole had good correlation with As exposure index in adult but in children. Correlation between skin dyspigmentation and As exposure index was observed in male adult. As exposure duration and skin symptoms also showed good correlation. Dyspigmentation of skin were prominent in male. Biological exposure monitoring showed weaker correlate to skin symptoms compare than past As exposure indexes. 2nd/1st methylation ratio of As was higher in children, then adult female than male. The results showed that gender difference and methylation potency have some role on status of skin lesions. Both hyperkeratosis and dyspigmentation were improved by water mitigation. Urine 8-OHdG, reflects DNA damage also decreased by avoiding As exposure. The intervention study showed that avoiding of As exposure was effective to the improvement of skin lesions and decrease the risk of carcinogenesis.

< See the slides on the next pages for the contents. >

Study of arsenism via consumption of arsenic contaminated ground water and intervention study in water mitigation in Inner Mongolia.

YOSHIDA, Takahiko
Department of Hygiene,
Asahikawa Medical College

1



2

Information of Gangfangying village

- One of 47 villages of Baotou city locates at the eastern boarder of city.
- Population 2080, representing 480 households.
- Major occupation, agriculture.
- Water for drinking and cooking:
 - In the past, six public open wells (shallow well).
 - As concentration; 0.00068ppm (only existing open well)
 - From 1979, gradually replaced by pump tube-well (deep well, approximately 330 wells)
 - Average As Conc.; 0.133ppm (ranging from 0.0001-1.79)
 - Pipeline water established at late August, 1999.
 - As concentration; 0.037ppm. All villagers were asked to use it.
- Arsenism
 - Appearance of first case; 1996.
 - Estimated As-related patients; 20%.
 - Short history of As exposure. No malignancy case.

3

Aim of study


- Assess the extent of As exposure until summer of 1999.
 - As exposure index: Σ (Conc. of well water x used years)
- Evaluate the severity of the symptoms with the correlation of individual As exposure.
 - Skin, cardio vascular, peripheral nerves, etc.
 - As exposure index, As exposure duration
- Research the factors that influence the symptoms.
 - Methylation of As, 8-OHdG (reflect DNA damage), etc.
- Evaluate potential changes in existing health effects by the intervention.
 - Intervention: change water source to lower As concentration.
 - Periodical follow-up health check after the change of water source.
 - July, 1999 as before; March, 2000 and September, 2000 as 6 or 12 months after.

4




Representative public open well in the village.
Only the existing open well in the village; North open well.

5




Representative personal pump-well in the villager's yard.


6




7




8




9




10




11




12




1 3




1 4




1 5




1 6




1 7




1 8




1 9




2 0




2 1




2 2




2 3




2 4



2 5



2 6



2 7




2 8



2 9




3 0




3 1




3 2



3 3



3 4




3 5



3 6



3 7




3 8



3 9



4 0



4 1

- ## Results
- Villagers had been exposed and were exposed to As.
 - Arsenic skin lesion:
 - Sever in male adult, especially at dyspigmentation
 - Arsenic exposure index:
 - Good correlate with hyperkeratosis in adult but in children.
 - Good correlate with dyspigmentation in male adult but in female adult and children.
 - Biological exposure monitoring:
 - Reflect the present As exposure.
 - Not reflect the symptoms as result of past exposure.
 - Symptom differences:
 - Gender • Age • Methylation ability

4 2

Results (continue)

- Influences of water source change
 - As exposure were reduced.
 - Skin alternations, major arsenism symptom were improved generally.
 - DNA damage were decreased.
- Improvement of water quality is effective to improve arsenism at least skin legions.

4 3

Recommendations

- Mixed exposure and health effects of other elements with As should be consider.
- Further follow-up should be conducted to find further health problems such as cancer even if their As exposure had stoped.
- Effectiveness for improvement of water quality should be evaluate on other health alternations.
 - Cardiovascular, peripheral nerves, pscology, etc.
- Epidemiological and experimental studies should be conducted for treatment and prevention of arsenism.
 - Study for symptom differences between population.
 - Gender, age, neutrition, another elements, etc.

4 4

Recommendations (Continue)

- Clinical surveys of residents and chemical analysis of water should be conducted in whole country wide.
- Improvement of water quality is necessary to stop occurring of new arsenism patients.
- Education
 - For residents
 - For government?

4 5

Session 3

Concluding Remarks

Major Points Confirmed and Recommendations for Future Studies

Michinori Kabuto

National Institute for Environmental Studies

Throughout this workshop, the current information on the affected areas with arsenic (and also fluoride in some parts) and populations in Bangladesh, West Bengal, Thailand and China has been overviewed and summarized, showing that population who have been drinking the underground water polluted with arsenic exceeding the international standards may count more than 100 millions in these areas. Although the skin lesions or symptoms such as hyper- or hypo-pigmentations or keratosis are most prevalent, it is expected that cases of various cancers including skin, lung etc. would increase if As measurements and disease surveillance are advanced. Moreover, if their exposures are prolonged among the people in highly polluted areas would continue chronically, an increasing trend of those diseases are predicted to be intensified and extended.

According to the several studies in Bangladesh and China, arsenic concentrations vary area-to-area and also even well-to-well in a small village, suggesting that much efforts are needed to diagnose each of the wells for good risk managements. And when high concentration is found for each of wells, however, it must be urgent to apply one of the most convenient methods or to change the water source to an alternative one in order to avoid its serious risks. With risk management, it is highly recommended to adopt the top-down type of governmental actions, since the affected areas are basically very poor in terms of economic status, risk knowledge as well as risk awareness and perception.

In the workshop, although the causal relationships between arsenic exposures and their specific effects have been indicated generally, further discussions especially with respect to recommendations for the future studies were also made based on those given by Dr. T. Yoshida, the last speaker, as shown in his slides. That is,

- 1) exposures and their health effects of other elements in addition to arsenic should be considered.
- 2) follow-up should be conducted to investigate more adverse effects such as cancer, since they may be expected even after exposures to arsenic are stopped.
- 3) effectiveness for improvement of water quality should be evaluate on other health alternations such as cardiovascular, peripheral nerves, psychology, etc.

- 4) epidemiological and experimental studies should be extended for prevention and treatment of arsenism with special attentions to symptom differences by population, gender, age, nutritional status, another elements etc.
- 5) clinical surveys on residents and chemical analysis of water should be extended to the whole country level.
- 6) improvement of water quality must be essential to prevent new cases of arsenism, and
- 7) education of appropriate knowledge for the residents as well as governors seems also essential.

Participant List

(as of March 2001)

議長 Chair person

兜 真徳 Michinori Kabuto

国立環境研究所 地域環境研究グループ

上席研究官

Director

Environment Risk Research Division, NIES

〒305-8506 つくば市小野川 16-2

16-2 Onogawa, Tsukuba, Ibaraki, 305-8506 Japan

Tel: +81-298-50-2333 Fax: +81-298-50-2571

e-mail: kabuto@nies.go.jp

大井 玄 Gen Ohi

国立環境研究所 所長

Director General, NIES

〒305-8506 つくば市小野川 16-2

16-2 Onogawa, Tsukuba, Ibaraki, 305-8506 Japan

Tel: +81-298-50-2300 Fax: +81-298-50-2854

e-mail: ohigen@nies.go.jp

吉田 貴彦 Takahiko Yoshida

旭川医科大学 医学部衛生学 教授

Professor

Department of Hygiene, Asahikawa Medical College

〒078-8510 旭川市緑が丘東 2 条 1-1

1-1 Higashi Nijo, Midorigaoka, Asahikawa,

Hokkaido, 078-8510 Japan

Tel: +81-166-68-2400/2402 Fax: +81-166-68-2409

e-mail: tyoshida@asahikawa-med.ac.jp

Amjad Shraim

Regional Environment Division NIES

〒305-8506 つくば市小野川 16-2

16-2 Onogawa, Tsukuba, Ibaraki, 305-8506 Japan

E-mail: amjad.shraim@nies.go.jp

渡辺 知保 Chiho Watanabe

東京大学医学部人類生態学教室 助教授

Associate Professor

Human Ecology, Faculty of Medicine

University of Tokyo

〒113-0033 文京区本郷 7-3-1

7-3-1 Hongou, Bunkyo, Tokyo, 113-0033 Japan

Tel: +81-3-5841-3487 Fax: +81-3-5841-3395

e-mail: chiho@humeco.mu-tokyo.ac.jp

金 銀龍 Jin Yinlong

中国予防医学科学院環境衛生及び衛生工程研究所
所長

Director

Institute of Environmental Health and Engineering,

Chinese Academy of Preventive Medicine

中国北京宣武区南緯路 29 路

29 Nan Wei Road Xuan Wu District Beijing 100050,
China

Tel: +86-10-6317-5273 Fax: +86-1-6317-5273

e-mail: jiniehe@public.bta.net.cn

孫 貴範 Sun Guifan

中国医科大学公共衛生院 院長

Dean & professor, School of Public Health,

China Medical University

中国遼寧省瀋陽市和平区北二馬路 92 市

Shenyang 110001, P. R. China

Fax: +86-24-23261744 or +86-24-86257545

何 公理 He Gongli
中国予防医学科学院 環境衛生・衛生工程研究所
科学部主任
Institute of Environmental Health and
Engineering, Chinese Academy of Preventive
Medicine
中国北京宣武区南緯路 29 路
29 Nan Wei Road Xuan Wu District Beijing, 100050
China

李 友 Li You
中国予防医学科学院
環境衛生・衛生工程研究所 研究助手
Researcher Assistant
Institute of Environmental Health and
Engineering, Chinese Academy of Preventive
Medicine
中国北京宣武区南緯路 29 路
29 Nan Wei Road Xuan Wu District
Beijing,100050 China
Tel: +86-10-6303-5945 Fax: +86-10-6301-3987
e-mail: liyoubj163.com

押川 尚子 Shoko Oshikawa
Prince of Songkla University Faculty of
Medicine
c/o Epidemiology Unit, Faculty of
Medicine, Prince of Songkla University
Hat yai, Songkla, 90112 Thailand
Tel: +66-74-429900 ex 1165~7 (電話は呼び出しの
都合上、日本時間の 15 時~ 22 時の間に限りま
す。)
Fax: +66-74-212900 or +66-74-212903
e-mail: oshoko@hotmail.com

Dipankar Chakraborti
Director
School of Environmental Study, Jadavpur University
Jadavpur University, Calcutta-700 032, India
Tel: +91-33-4735233 Fax: +91-33-4734266
e-mail: dcsoesju@vsnl.com

遠山 千春 Chiharu Tohyama
国立環境研究所 環境保健部 部長
Director
Environmental Health Sciences Division NIES
〒305-8506 つくば市小野川 16-2
16-2 Onogawa, Tsukuba, Ibaraki,
305-8506 Japan
Tel: +81-298-50-2336 Fax: +81-298-50-2574
e-mail: ctohyama@nies.go.jp

平野 靖史郎 Seishiro Hirano
国立環境研究所 地域環境研究グループ
主任研究員
Independent Senior Researcher
Regional Environment Division, NIES
〒305-8506 つくば市小野川 16-2
16-2 Onogawa, Tsukuba, Ibaraki,
305-8506 Japan
Tel: +81-298-50-2512 Fax: +81-298-50-2512
e-mail: seishiro@nies.go.jp

曾根 秀子 Hideko Sone
国立環境研究所 地域環境研究グループ
主任研究員
Independent Senior Researcher
Regional Environment Division, NIES
〒305-8506 つくば市小野川 16-2
16-2 Onogawa, Tsukuba,
Ibaraki, 305-8506 Japan
Tel: +81-298-50-2464 Fax: +81-298-50-2571
e-mail: hsone@nies.go.jp

Jia Guang
国立環境研究所 STA フェロー
NIES STA Fellow

大野 浩一 Kohichi Ohno
北海道大学工学研究科
環境リスク工学分野 助手
Research Associate
Hokkaido University Faculty of Engineering
〒060-0808 札幌市北区北 8 条西 5 丁目
Kita8,Nishi5,Kitaku,Sapporo,060-0808 Japan
Tel & Fax: +81-11-706-7280
E-mail: ohnok@eng.hokudai.ac.jp

佐藤 裕子 Yuko Satoh
北海道大学 工学研究科
環境リスク工学分野 博士後期課程学生
Graduate Student
Hokkaido University Faculty of Engineering
〒060-0808 札幌市北区北 8 条西 5 丁目
Kita8,Nishi5, Kitaku, Sapporo,060-0808 Japan
Tel & Fax: +81-11-706-7280
E-mail: satoyuko@eng.hokudai.ac.jp

鈴木 和夫 Kazuo Suzuki
千葉大学薬学部 衛生化学研究室
Chiba University
Faculty of Pharmaceutical Sciences
〒263-8522 千葉市稲毛区弥生町 1-33
1-33 Yayoicho, Inageku, Chiba, 263-8522 Japan
Tel & Fax: +81-43-290-2891
E-mail ktsuzuki@p.chiba-u.ac.jp

Badal K. Mandal
千葉大学薬学部衛生化学研究室
JSPS Fellow
Chiba University
Faculty of Pharmaceutical Sciences
Te/Fax: +81-43-290-2892

浅見 真理 Mari Asami
国立公衆衛生院 水道工学部
National Institute of Public Health
Department of Water Supply Engineering
〒108-8638 東京都港区白金台 4-6-1
4-6-1 Shiroganedai, Minatoku, Tokyo, 108-8638 Japan
Tel: +81-3-3441-7111 Fax: +81-3-3446-4314
E-mail: mari@iph.go.jp

山本 玲子 Reiko Yamamoto
尚絢女学院短期大学 教授
Professor
Shokei Women's Junior College
〒981-1213 宮城県名取市ゆりが丘 4-10-1
4-10-1 Yurigaoka, Natorishi, Miyagi,981-1213 Japan
Tel & Fax: +81-22-383-0202
E-mail: yamamoto@shokei.ac.jp

山内 博 Hiroshi Yamauchi
聖マリアンナ医科大学予防医学教室 助教授
Associate Professor
St. Marianna University School of Medicine
Department of Preventive Medicine
〒216-8511 川崎市宮前区菅生 2-16-1
2-16-1 Sugao, Miyamaeku, Kawasakishi
Kanagawa 216-8511, Japan
Fax & Tel +81-44-977-0268
E-mail: h2yama@marianna-u.ac.jp

有蘭 幸司 Koji Arizono
熊本県立大学 環境共生学部 教授
Professor
Prefectural University of Kumamoto,
Faculty of Environmental and Symbiotic Sciences
〒862-8502 熊本市月出 3-1-100
3-1-100 Tsukide, Kumamoto, 862-8502 Japan
Tel& Fax: +81-96-383-6062
e-mail: arizono@pu-kumamoto.ac.jp

小久保順子 Junko Kokubo
群馬大学 保健学科看護学 助手
Research Associate
Gunma University School of Health Sciences
〒371-8511 前橋市昭和町 3-39-22
3-39-22 Showamachi, Maebashishi, Gunma
371-8511 Japan
Tel/Fax: +81-27-220-8983
E-mail: jkokubo@health.gunma-u.ac.jp

ザイ 国方 Zhai Guofang
ダイナボット株式会社 学術部
〒106-8535 東京都港区六本木 1-9-9
六本木ファーストビル
Roppongi First bld. 1-9-9 Roppongi, Minatoku,
Tokyo, 106-8535 Japan
E-mail: guofang.zhai@abbott.com

Yaxiong Xie
Juntendo University School of Medicine
Department of Epidemiology and
Environmental Health
2-1-1, Hongo, Bunkyo-ku Tokyo, 113-8421 Japan
Tel: +81-3-5802-1047 Fax: +81-3-3812-1026
E-mail: yaxxie@hotmail.com (Japanese
cannot be read at this e-mail address)

風間 駿 Shun Kazama
〒154-0002 東京都世田谷区下馬 3-31-16
3-31-16 Shimouma, Setagayaku, Tokyo,
154-0002 Japan
Tel: 03-3422-9611 Fax: 03-3422-9617
E-mail: kazama@dp.catv.ne.jp

大手
流通戦略総合研究所
e-mail: zbb23224@boat.zero.ad.jp

稲岡 司 Tsukasa Inaoka
熊本大学医学部 公衆衛生学教室 助教授
Associate Professor
School of Medicine, Kumamoto University
〒860-0811 熊本市本荘 2-2-1
2-2-1 Honjo, Kumamoto, 860-0811 Japan
Tel: +81-96-373-5111 Fax: +81-96-373-5113
e-mail: inaoka@gpo.kumamoto-u.ac.jp

山村 尊房 Sombo Yamamura
国連大学高等研究所
The United Nations University
Institute of Advanced Studies
〒150-8304 東京都渋谷区神宮前 5-53-67
5-53-67 Jingumae, Shibuya, Tokyo, 150-8304 Japan
Tel: +81-3-5467-2811 Fax: +81-3-5467-2324
e-mail: yamamura@ias.unu.edu

大塚 健司 Kenji Ohtsuka
日本貿易振興会アジア経済研究所開発研究部
Institute of Developing Economies JETRO
〒261-8545 千葉県美浜区若葉 3-2-2
3-2-2 Wakaba, Mihama, Chiba, 261-8545 Japan
Tel: +81-43-299-9752 (直通)
Fax: +81-43-299-9763 (研究棟 3F)
e-mail: otuka@ide.go.jp

辻田 祐子 Yuko Tsujita
日本貿易振興会アジア経済研究所地域研究第1部
Institute of Developing Economies JETRO
〒261-8545 千葉県美浜区若葉 3-2-2
3-2-2 Wakaba, Mihama, Chiba, 261-8545 Japan
Tel: +81-43-299-9605 Fax: +81-43-299-9729
e-mail: tsujita@ide.go.jp

Dr. Tarit Roychowdhury
NIHS
Mrs. Sanchita Roychowdhury
NIHS

河原 純子 Junko Kawahara
東京大学大学院修士課程新領域創成科学研究科
環境学専攻
Tokyo University, Graduate School of Frontier Sciences
〒113-0033 東京都文京区本郷 7-3-1
7-3-1 Hongo, Bunkyo, Tokyo, 113-0033 Japan,
Tel: +81-3-5841-7335
e-mail: kawahara@yy.t.u-tokyo.ac.jp

Zafar ADEEL
Academic Programme Officer
The United Nations University Environment
and Sustainable Development
〒150-8925 渋谷区神宮前 5-53-70
5-53-70 Jingumae, Shibuyaku, Tokyo, 150-8925 Japan
Tel: + 81-3-3499-2811 Fax: +81-3-3406-7347
e-mail: Adeel@hq.unu.edu
URL: <http://www.unu.edu/env>

高橋 徹 Tohru Takahashi
熊本県立大学環境共生学部 海洋生態学研
究室 (堤研)
Minamata Environmental Research
& Develop Co., Ltd. Visiting researcher,
Active Collaboration Center for
Environmental & Symbiotic Sciences
Prefectural University of Kumamoto
〒862-8502 熊本市月出 3-1-100
3-1-100 Tsukide, Kumamoto, 862-8502 Japan
Tel: +81-96-383-2929 内454 Fax: +81-096-384-6765
e-mail: tohru@pu-kumamoto.ac.jp
rhizo@poem.ocn.ne.jp

篠原 直秀 Naohide Shinohara
東京大学大学院新領域創成科学研究科 修士2年
Tokyo University, Graduate School of Frontier Sciences
〒113-0033 東京都文京区本郷 7-3-1
7-3-1 Hongo, Bunkyo, Tokyo, 113-0033 Japan
e-mail: shinohara@yy.t.u-tokyo.ac.jp

小椋 康光 Yasumitsu Ogura
千葉大学薬学部 衛生化学研究室 助手
Research Associate Chiba University
Faculty of Pharmaceutical Sciences
〒263-8522 千葉市稲毛区弥生町 1-33
1-33 Yayoicho, Inageku, Chiba, 263-8522 Japan
Tel & Fax: +81-43-290-2892
e-mail: ogra@p.chiba-u.ac.jp

富田 貴之 Takayuki Tomita
千葉大学大学院衛生化学研究室 修士1年
Chiba University, Graduate School of Medicine
〒263-8522 千葉市稲毛区弥生町 1-33
1-33 Yayoicho, Inageku, Chiba, 263-8522 Japan
Te/Fax: +81-43-290-2892
e-mail: tomy@p.chiba-u.ac.jp
〒260-0044 千葉県千葉市中央区松波 3-5-10
SK 松波 102
Tel: +81-43-287-5212

徳永 裕司 Hiroshi Tokunaga
国立医薬品食品衛生研究所
National Institute of Health Sciences
〒158-8501 東京都世田谷区上用賀 1-18-
1-18-1 Kamiyoga, Setagayaku, Tokyo, 158-8501 Japan
Tel: +81-3-3700-1141 Fax: +81-3-3707-6950
e-mail: tokunaga@nihs.go.jp

山本 和夫 Kazuo Yamamoto
東京大学環境安全研究センター 教授
Professor
Environmental Science Center, University of Tokyo
〒113-0033 東京都文京区本郷 7-3-1
7-3-1 Hongo, Bunkyo, Tokyo, 113-0033 Japan
Tel: +81-3-5841-2972, +81-3-5841-2973 (事務室)
Fax: +81-3-3813-7294

宮崎 賀織 Kaori Miyazaki

東京大学大学院医学系研究科

人類生態学教室

University of Tokyo, Graduate School of Medicine

〒113-0033 文京区本郷 7-3-1

7-3-1 Hongo, Bunkyo, Tokyo, 113-0033 Japan

Tel: +81-3-5841-3529 Fax: +81-3-5841-3395

e-mail: kaori@humeco.m.u-tokyo.ac.jp

崔星

東京大学 薬学部遺伝学教室

University of Tokyo

Faculty of Pharmaceutical Sciences

〒113-0033 文京区本郷 7-3-1

7-3-1 Hongo, Bunkyo, Tokyo, 113-0033 Japan

REPORT RESEARCH FROM
THE NATIONAL INSTITUTE FOR ENVIRONMENTAL STUDIES, JAPAN

No.166

国立環境研究所研究報告 第 166 号
(R - 166 - 2001)

This Proceeding is available by request from :
Michinari Kabuto, DHSc Acting Director for Health Researches

Tel : +81-298-50-2333

Fax : +81-298-50-2571

e-mail : kabuto@nies.go.jp

【平成 13 年 5 月 31 日編集委員会受付】

【平成 13 年 7 月 4 日編集委員会受理】

平成 13 年 8 月 31 日発行

発 行 独立行政法人 国立環境研究所

〒305-8506 茨城県つくば市小野川 16 番 2

電話 0298 - 50 - 2343 (ダイヤルイン)

印 刷 前田印刷株式会社 筑波支店

〒305-0033 茨城県つくば市東新井 14 - 3

Published by the National Institute for Environmental Studies
16-2 Onogawa, Tsukuba, Ibaraki 305-8506 Japan
August 2001

本報告書は再生紙を使用しています。