





# The OCO-2 and OCO-3 missions: Status, results and plans


Vivienne Payne, Abhishek Chatterjee, Junjie Liu Jet Propulsion Laboratory, California Institute of Technology

And the OCO-2/OCO-3 Project Teams, Science Team!



### Tracking the whole-atmosphere CO<sub>2</sub> growth rate in near real time






CO<sub>2</sub> growth rate for 2024: One of the highest on record

(Both for NOAA ground-based record and for satellite record)

Strongly decoupled from ENSO index, likely that anomaly originated in the Southern Tropics

- Pandey et al. (2024)
  AGU Advances,
  10.1029/2023AV00114
  5
- Pandey et al. (in preparation)



### The OCO-2 and OCO-3 missions




### **Orbiting Carbon Observatory-2 (OCO-2)**

- Launched July 2, 2014
- Free flying spacecraft
- Sun-synchronous polar orbit



#### **Orbiting Carbon Observatory-3 (OCO-3)**

- Launched May 4, 2019
- International Space Station (ISS)
- ± 52° inclined orbit



OCO-2 and OCO-3 data products are publicly available via the NASA GES DISC


### OCO-2 and OCO-3: Complementary coverage and sampling

#### OCO-2

- "Pole-to-pole" coverage, depending on season;
- Fixed 1330h equator crossing time (and local overpass time)

#### **OCO-3**

- Coverage limited to ±52° latitude, changing with season;
- Observations span all times of day




### OCO-2 and OCO-3: Complementary coverage and sampling

#### OCO-2

- "Pole-to-pole" coverage, depending on season;
- Fixed 1330h equator crossing time (and local overpass time)

#### **OCO-3**


- Coverage limited to ±52° latitude, changing with season;
- Observations span all times of day





# OCO-3 Snapshot Area Maps (SAMs)


- Collects data over an ~80 km × 80 km area in 2 minutes
- Complements the near-global nadir & glint measurements




See R. Nelson/A. Chatterjee talk on Tuesday!



# OCO-2: Continuing to age like a fine wine!



- Instrument and spacecraft are in excellent health
- Outgassing/icing effects continue to slow as the mission extends in time




Figures: R. Rosenberg and the OCO-2 dal team




# OCO-3: With age (and v11) comes wisdom

- OCO-3 instrument is in excellent health!
- OCO-3 v11: Consistency with OCO-2 v11.2 L2 algorithm, plus OCO-3-specific improvements to L1B calibration and geolocation compared to v10










# Level 2 XCO<sub>2</sub> and SIF products

- Latest and greatest Level 2 data versions: v11.2 for OCO-2, v11 for OCO-3
  - Consistent algorithm used for OCO-2 and OCO-3!

Figure: Chris O'Dell, CSU

#### Forward stream


- Subset of soundings, latency < 1 week</li>
- Products removed from GES DISC site on rolling basis
- New! Bias-corrected forward stream L2 products
  - Intended for users who need low latency



OCO-2 coverage, forward stream (06/22)

### Retrospective stream

- All "cloud-free" soundings processed, monthly cadence.
- Full record available at GES DISC site
- L2 Lite products: Bias correction, quality filter applied
  - · Recommended for most users




OCO-2 coverage, retrospective stream (06/22)



# Why "OCO-2 v11.2" but "OCO-3 v11"?


- OCO-2 v11 → v11.1: Switch to Copernicus Digital Elevation Map
- OCO-2 v11.1 → v11.2: Transition from GEOS-FPIT to GEOS-IT met fields
  - Date of GEOS-IT transition: 2<sup>nd</sup> April 2024
  - Users should be aware: Transition introduces subtle but non-zero discontinuity in record
    - Mitigated, but not fully addressed, by v11.2 Lite reprocessing for pre-2024 data
    - To fully address this, a full L2FP reprocessing would be required
  - For more info, see OCO-2 Data User Guide



- OCO-3 v11 is fully consistent throughout the record
  - OCO-3 v11 processing started later, was able to "bake in" these DEM, GEOS-IT updates




### Validation against TCCON (GGG2020): Agreement < 1 ppm



# TCCON GGG2020.1 status




Changes to Xgases from implementation of time-varying  $O_2$  DMF



#### Progress:

- Consistent time-varying O<sub>2</sub> DMF for all Xgases
- N<sub>2</sub>O T-dependent bias reduction
- Track use of alternate CO priors
- $\Sigma$  Upgrade public file creation to support above
- $\Sigma$  Generate alternate pre-Apr 2024 CO priors

### Changes to XN<sub>2</sub>O from implementation of T-dependent bias reduction




- Impact of new time-varying  $O_2$  DMF is much smaller than the TCCON error budget, benefit is to long timeseries analyses
- XN<sub>2</sub>O bias reduction reduces temperature dependence in the common temperature ranges, future work needed to address the coldest temperatures
- Updating TCCON infrastructure to ensure that the GGG2020 to GGG2020.1 processing is applied correctly is taking longer than expected
  - First of three components almost complete
- Aiming now to have development complete by Sep. 2025.
- Sites that wish to use improved CO priors will need time after the GGG2020.1 release to reprocess



# "Can I use OCO-2 and OCO-3 together?"

### Yes!

- Direct comparisons for XCO2:
  - OCO-3 v11 agrees with OCO-2 v11.2 to better than 0.5 ppm for co-located soundings
  - This is well within the agreement of either sensor with TCCON.
  - Taylor et al. (2025), in prep





### Closing remarks

- OCO-2 and OCO-3 are both in excellent health
- With OCO-2 v11.2 and OCO-3 v11, L2 products are now in step
- Accuracy and precision of XCO<sub>2</sub> products:
  - < 1 ppm for **both missions**
- Characterization of a new instrument/data product can take some time....
- XCO<sub>2</sub> and SIF datasets from OCO-2 and OCO-3
  - Shining new light on natural and anthropogenic components of carbon cycle
- Length of the record and quality of the data products are key
  - Particularly for studies of the natural carbon cycle